## **Attention to prevention**



Your Guide to Preventive Care for Baby's 1st and 2nd Year

When it comes to patient care HPSJ is on your team. We understand that preventive health care is about improving quality of life. This quick reference guide is here to help you reach those goals with your patients.

At all visits

If there is risk (or if immunization was not given prior) 7-20 years 5 month 24 month 18 month 2 month Newborn 11 years month month month month 14 years 5 years month 10 years 2 years 13 years 2-4 day 3 years 6 years 4 years 6 years 8 years 5 years  $\sim$ 9 0 Well Child Visits Head Measurement Height, Weight, Blood Pressure Disease Screening Lead Screening Blood test (anemia) **Cholesterol Screening** Urinalysis TB test Vision & Hearing Screening Referral to Dentist Fluoride Varnish Application Development, Behavior, Past Safety, Nutrition, Parenting Tobacco assessment Anticipatory Guidance **CRAFFT** Assessment **Depression Screening** STD Screening and Education **Immunizations** Hepatitis B **Rotovirus** Tetanus, Diptheria, Pertussis (Tdap) Haemophilus influenza type B (Hib) Pneumococcal (PCV) Inactivated Polio Vaccine (IPV) Measles, Mumps, Rubella (MMR) Varicella (Chickenpox) Hepatitis A Influenza Human Papillomavirus (HPV)\* Meningococcal Conjugate

Source: CDC Immunization Schedules for all Ages. Periodicity Schedule, American Academy of Pediatrics. This material can be provided in larger font and/or other languages. Please call (888) 936- PLAN (7526) \*All 11 or 12 year olds – both girls and boys – should receive 3 doses of HPV vaccine to protect against HPV-related disease. The full HPV vaccine series should be given as recommended for best protection.