

# MEDICATION COVERAGE POLICY



## PHARMACY AND THERAPEUTICS ADVISORY COMMITTEE

<b>POLICY:</b>	Ankylosing Spondylitis (AS)	<b>P&amp;T DATE:</b>	05/14/2019
<b>CLASS:</b>	Rheumatology/Anti-inflammatory Disorders	<b>REVIEW HISTORY</b>	02/08, 05/10, 02/12,
<b>LOB:</b>	Medi-Cal	(month/year)	10/14, 02/16, 02/17, 02/18

*This policy has been developed through review of medical literature, consideration of medical necessity, generally accepted medical practice standards, and approved by the HPSJ Pharmacy and Therapeutic Advisory Committee.*

## OVERVIEW

Ankylosing Spondylitis (AS) is an inflammatory condition that usually involves the spine.<sup>1</sup> Unlike rheumatoid arthritis (RA), oral DMARDs (methotrexate, leflunomide, etc) have not been effective in the treatment of AS. NSAIDs (ibuprofen, naproxen, etc) and physical therapy are first-line treatment. In patients who are symptomatic despite NSAID treatment, treatment with TNF biologics are recommended. This review will examine the treatment guidelines of AS, the currently available AS drug products, and their coverage criteria.

**Table 1. Available Ankylosing Spondylitis Agents (Current as of 2/2019)**

Formulary Agents					
Therapeutic Class	Generic Name (Brand Name)	Available Strengths	Fml Limits	Cost/ Month	Notes
Biologics	Adalimumab (Humira, Humira CF)	20mg/0.4ml, 40mg/0.8ml	PA; PL; SP	\$5,371.99	Reserved for treatment failure to 2 NSAIDs tried within the last 30-60 days.
	Etanercept (Enbrel)	50mg/ml, 25mg/ml,	PA; PL; SP	\$4,948.98	Reserved for treatment failure to 2 NSAIDs tried within the last 30-60 days.
	Infliximab-dyyb (Inflectra) Infliximab-abda (Renflexis)	100mg IV vial	PA; PL; SP	--	Reserved for treatment failure to 2 NSAIDs tried within the last 30-60 days.
	Infliximab (Remicade)	100mg IV vial	NF	--	Non-Formulary
	Golimumab (Simponi)	50mg/0.5ml, 100mg/ml	PA; PL; SP	\$4,610.15	Reserved for treatment failure to either Adalimumab, Etanercept, or Infliximab.
	Certolizumab (Cimzia)	200mg/ml	PA; PL; SP	--	Reserved for treatment failure to either Adalimumab, Etanercept, or Infliximab.
	Secukinumab (Cosentyx)	150mg/ml	PA; PL; SP	\$5,261.11	Reserve for treatment failure to two 1st line agents OR one 1st line agent and one 2nd line agent

PL = Prescriber Limit, (Must be prescribed by Rheumatologist); PA = Prior Authorization Required; NF = Non-Formulary; SP = Specialty

## ⊕ EVALUATION CRITERIA FOR APPROVAL/EXCEPTION CONSIDERATION

Below are the coverage criteria and required information for each agent. These coverage criteria have been reviewed approved by the HPSJ Pharmacy & Therapeutics (P&T) Advisory Committee. For conditions not covered under this Coverage Policy, HPSJ will make the determination based on Medical Necessity as described in HPSJ Medical Review Guidelines (UM06).

### Biologics

#### **1<sup>st</sup> line—Adalimumab (Humira, Humira CF), Etanercept (Enbrel), Infliximab (Infliximab, Remicade)**

- Coverage Criteria:** Reserved for documented symptomatic AS despite treatment with NSAIDs (unless NSAID-intolerant). An adequate trial is defined as at least 2 different NSAIDs tried over 1 month or 2 different NSAIDs over 2 months.
- Limits:** None
- Required Information for Approval:** Prescription history showing at least 2 NSAIDs tried.
- Other Notes:** Medication is to be dispensed by HPSJ's designated specialty pharmacy. Must be initiated by a rheumatologist. Biologics exceeding labeled standard maintenance doses may be approved 1 month at a time. Subsequent fills of the increased maintenance dose will require documentation of symptom improvement.
- Non Formulary:** Remicade

#### **2<sup>nd</sup> line—Certolizumab (Cimzia), Golimumab (Simponi)**

- Coverage Criteria:** Reserved for treatment failure to Adalimumab, Etanercept, or Infliximab.
- Limits:** None
- Required Information for Approval:** Prescription history showing at least 3 month trial of one first line agent (Adalimumab, Etanercept, or Infliximab).
- Other Notes:** Medication is to be dispensed by HPSJ's designated specialty pharmacy. Must be initiated by a rheumatologist. Biologics exceeding labeled standard maintenance doses may be approved 1 month at a time. Subsequent fills of the increased maintenance dose will require documentation of symptom improvement.

#### **3<sup>rd</sup> line—Secukinumab (Cosentyx)**

- Coverage Criteria:** Reserved for treatment failure/documentated intolerance to two 1<sup>st</sup> line agents (adalimumab, etanercept, infliximab) OR one 1<sup>st</sup> line agent (adalimumab, etanercept, infliximab) and one 2<sup>nd</sup> line agent (certolizumab, golimumab). Must be prescribed by rheumatologist.
- Limits:** None
- Required Information for Approval:** Prescription history showing at least 3 month trial of [1] adalimumab, etanercept, or infliximab AND/OR [2] certolizumab or golimumab.
- Other Notes:** Medication is to be dispensed by HPSJ's designated specialty pharmacy. Must be initiated by a rheumatologist. Biologics exceeding labeled standard maintenance doses may be approved 1 month at a time. Subsequent fills of the increased maintenance dose will require documentation of symptom improvement.


## CLINICAL JUSTIFICATION

The goals of treatment are to reduce symptoms to maintain body function and quality of life. The 2015 American College of Rheumatology (ACR)/Spondylitis Association of America (SAA)/Spondyloarthritis Research and Treatment Network (SRTN) Guidelines<sup>2</sup> recommends the following:

### Active AS

- NSAIDs and physical therapy are first-line treatment.
  - The guidelines define “adequate trial” as “lack of response (or intolerance) to at least 2 different NSAIDs over 1 month or incomplete responses to at least 2 different NSAIDs over 2 months.”
- In patients who are symptomatic despite NSAID treatment, treatment with TNF biologics are recommended.
  - There is insufficient evidence to favor one TNF biologic over another. However, experts agreed that in patients with AS and inflammatory bowel disease, infliximab or adalimumab is preferred over etanercept due to lower rates of iritis.
  - For patients with active AS despite treatment with TNF biologic, the guidelines recommend switching to another TNF biologic (as opposed to adding a DMARD).
- Methotrexate and Leflunomide have shown to have minimal benefit and are associated with side effects. The benefits did not outweigh the risks and, therefore, are generally, not recommended.
  - Sulfasalazine was shown to have a small benefit on pain relief and may be an option for patients who cannot use TNF biologics.
  - DMARDs are preferred over non-TNF biologics (abatacept, tocilizumab, ustekinumab, etc) due to questionable efficacy and study bias.
- Systemic glucocorticoids are not recommended due to lack of strong safety and efficacy data.

Figure 1: ACR/SAA/SRTN Active AS Treatment Algorithm




### Stable AS

- Patients with stable AS or on stable treatment regimen, experts recommend using NSAIDs on an as-needed basis.
- Patients with stable AS receiving both a TNF biologic and NSAIDs or a TNF biologic with DMARDs, may consider discontinuing the NSAID or DMARD and continuing on TNF biologic as monotherapy.

The efficacy between TNF biologics do not differ significantly but the cost may vary due to differences in administration frequency (twice monthly vs. weekly vs. monthly, and so forth). Therefore, HPSJ’s order of preference of the biologic therapies are based on the cost-benefit ratio where the first-line biologics are agents associated with the lowest cost-benefit ratio.

Figure 2: ACR/SAA/SRTN Stable AS Treatment Algorithm



## ☒ REFERENCES

1. Ruderman E and Tambar S. Rheumatoid Arthritis. American College of Rheumatology. Updated August 2012. Accessed on June 22, 2015. Available at: [https://www.rheumatology.org/Practice/Clinical/Patients/Diseases\\_And\\_Conditions/Rheumatoid\\_Arthritis/](https://www.rheumatology.org/Practice/Clinical/Patients/Diseases_And_Conditions/Rheumatoid_Arthritis/)
2. Singh J, Furst DE, Bharat A, et al. 2012 Update of the 2008 American College of Rheumatology Recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *American College of Rheumatology*. 2012; 64(5): 625-639.
3. Yoo DH, Racewicz A, Brzezicki J, et al. A phase III randomized study to evaluate the efficacy and safety of CT-P13 compared with reference infliximab in patients with active rheumatoid arthritis: 54-week results from the PLANETRA study. *Arthritis Res Ther* 2016;18:82.
4. ClinicalTrials.Gov. Efficacy and Safety Study of ABP 501 Compared to Adalimumab in Subjects With Moderate to Severe Rheumatoid Arthritis. October 23, 2016. <https://clinicaltrials.gov/ct2/show/NCT01970475>. Accessed January 22, 2017.
5. Griffiths CE, et al. The EGALITY study: A confirmatory, randomised, double-blind study comparing the efficacy, safety and immunogenicity of GP2015, a proposed etanercept biosimilar, versus the originator product in patients with moderate to severe chronic plaque-type psoriasis. *Br J Dermatol*. 2016 Oct 27. doi: 10.1111/bjd.15152.
6. Dapavo P, et al. The infliximab biosimilar in the treatment of moderate to severe plaque psoriasis. *Journal of American Academy of Dermatology*. 2016 Oct;75(4):736-9.
7. FIMEA. Interchangeability of Biosimilars—Position of Finnish Medicines Agency Fimea. May 22, 2015. [http://www.fimea.fi/instancedata/prime\\_product\\_julkaisu/fimea/embeds/fimeawwwstructure/29197\\_Biosimilaarien\\_vaihtokelpoisuus\\_EN.pdf](http://www.fimea.fi/instancedata/prime_product_julkaisu/fimea/embeds/fimeawwwstructure/29197_Biosimilaarien_vaihtokelpoisuus_EN.pdf). Accessed January 22, 2017.
8. FDA. Summary Minutes of the Arthritis Advisory Committee Meeting. July 12, 2016. <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/ArthritisAdvisoryCommittee/UCM520027.pdf>. Accessed January 21, 2017.
9. Dörner T, Strand V, Cornes P, et al. The changing landscape of biosimilars in rheumatology. *Ann Rheum Dis* 2016; 75:974–82. doi:10.1136/annrheumdis-2016-209166
10. FDA. Biosimilars: Questions and Answers Regarding implementation of Biologics Price Competition and Innovation Act of 2009. April 2015. <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM444661.pdf>. Accessed January 21, 2017
11. Dapavo P, et al. The infliximab biosimilar in the treatment of moderate to severe plaque psoriasis. *Journal of American Academy of Dermatology*. 2016 Oct;75(4):736-9
12. Results from the NOR-SWITCH study support switch from Remicade to Remsima (biosimilar infliximab). Mundipharma. 19 October 2016. <http://www.mundipharma.com/docs/default-source/default-document-library/161019-ueg-press-release-final.pdf?sfvrsn=0>. Accessed 2 Feb 2017.
13. Baeten D, Sieper J, Braun J, et al. Secukinumab, an Interleukin-17A Inhibitor, in Ankylosing Spondylitis. *N Engl J Med*. 2015;373(26):2534-48.
14. Maxwell LJ, Zochling J, Boonen A, et al. Anti-TNF-alpha drugs for treating ankylosing spondylitis. *Cochrane Review*. Last updated 18 April 2015. Accessed 6 February 2016. Available at: [http://www.cochrane.org/CD005468/MUSKEL\\_anti-tnf-alpha-drugs-for-treating-ankylosing-spondylitis](http://www.cochrane.org/CD005468/MUSKEL_anti-tnf-alpha-drugs-for-treating-ankylosing-spondylitis).
15. Maxwell LJ, Zochling J, Boonen A, et al. Anti-TNF-alpha drugs for treating ankylosing spondylitis. *Cochrane Review*. Last updated 18 April 2015. Accessed 6 February 2016. Available at: [http://www.cochrane.org/CD005468/MUSKEL\\_anti-tnf-alpha-drugs-for-treating-ankylosing-spondylitis](http://www.cochrane.org/CD005468/MUSKEL_anti-tnf-alpha-drugs-for-treating-ankylosing-spondylitis).
16. Landewe R, Braun J, Deodhar A, et al. Efficacy of certolizumab pegol on signs and symptoms of axial spondyloarthritis including ankylosing spondylitis: 24-week results of a double-blind randomized placebo-controlled Phase 3 study. *Ann Rheum Dis*. 2014; 73(1): 39-47.

17. Van der Heijde D, Ramiro S, Landewé R, et al 2016 update of the ASAS-EULAR management recommendations for axial spondyloarthritis *Annals of the Rheumatic Diseases* 2017;76:978-991.
18. Kay et al. Consensus-based recommendations for the use of biosimilars to treat rheumatological diseases. *Ann Rheum Dis* 2018;77:165–174.
19. Tanaka Y, Yamanaka H, Takeuchi T, Inoue M, Saito K, Saeki Y, et al. Safety and efficacy of CT-P13 in Japanese patients with rheumatoid arthritis in an extension phase or after switching from infliximab. *Modern Rheumatol* 2017;27:237–45.
20. Park W, Yoo DH, Miranda P, Brzosko M, Wiland P, Gutierrez-Urena S, et al. Efficacy and safety of switching from reference infliximab to CT-P13 compared with maintenance of CT-P13 in ankylosing spondylitis: 102-week data from the PLANETAS extension study. *Ann Rheum Dis* 2017;76:346–54.
21. Jorgensen KK, Olsen IC, Goll GL, Lorentzen M, Bolstad N, Haavardsholm EA, et al. Switching from originator infliximab to biosimilar CT-P13 compared with maintained treatment with originator infliximab (NOR-SWITCH): a 52-week, randomised, double-blind, non-inferiority trial. *Lancet* 2017;389:2304–16.
22. Yoo DH, Prodanovic N, Jaworski J, Miranda P, Ramiterre E, Lanzon A, et al. Efficacy and safety of CT-P13 (biosimilar infliximab) in patients with rheumatoid arthritis: comparison between switching from reference infliximab to CT-P13 and continuing CT-P13 in the PLANETRA extension study. *Ann Rheum Dis* 2017;76:355–63.
23. Bridges, S. L., White, D. W., Worthing, A. B., Gravallese, E. M., O'Dell, J. R., Nola, K., Kay, J., Cohen, S. B. and on behalf of the American College of Rheumatology (2018), The Science Behind Biosimilars. *Arthritis Rheumatol*. doi:10.1002/art.40388

## **REVIEW & EDIT HISTORY**

Document Changes	Reference	Date	P&T Chairman
Creation of Policy	Biological Response Modifiers Review 2-19-08.doc	2/2008	Allen Shek, PharmD
Updated Policy	Biologic Response Modifiers 2010 final.docx	5/2010	Allen Shek, PharmD
Updated Policy	TNF MUE summary 2-21-2012.docx	2/2012	Allen Shek, PharmD
Updated Policy	Psoriatic Arthritis & Ankylosing Spondylitis.docx	10/2014	Jonathan Szkotak, PharmD
Updated Policy	Class Review- Biologics, Apremilast, and Tofacitinib in Inflammatory Joint, Skin, and Bowel Diseases.docx	2/2016	Johnathan Yeh, PharmD
Updated Policy	Class Review- Biologics, Apremilast, and Tofacitinib in Inflammatory Joint, Skin, and Bowel Diseases.docx	02/2017	Johnathan Yeh, PharmD
Updated Policy	HPSJ Coverage Policy – Rheumatology – Ankylosing Spondylitis 2018-02.docx	02/2018	Johnathan, Yeh, PharmD
Updated Policy	HPSJ Coverage Policy – Rheum & Immuno – Ankylosing Spondylitis 2019-05.docx	05/2019	Matthew Garrett, PharmD

*Note: All changes are approved by the HPSJ P&T Committee before incorporation into the utilization policy*