

Vaccination Schedule for Children

Well-Child Check-up & Shots: Children from birth to 17 years old

The Centers for Disease Control and Prevention (CDC) recommends annual flu vaccines for everyone age 6 months or older. Check with your doctor when flu season starts to see if it's right for your child. At your child's next visit, make sure to tell your doctor if your child will be traveling out of the US, they may need extra shots. If your child is behind on shots, your doctor can make a plan to get all their shots done.

 Well-Child Check-up  Blood Screen  Shots  Vision Screen  Hearing Screen  Ask about the flu shot	
Age	Recommended Vaccines & Tests
Birth	   (Hepatitis B)
3-5 days	
7-14 days	
2 months	  (DTaP, Hep B, Hib, PCV, Rotavirus, IPV)
4 months	  (DTaP, Hib, PCV, Rotavirus, IPV)
6 months	  (DTaP, Hep B, PCV, Hib & Rotavirus if needed, IPV) 
9 Months	 
12 months	  (MMR, Hepatitis A, Varicella, Hib if needed, PCV) 
15-18 months	  (DTap) 
2 years	  (Hepatitis A) 
3 years	 
4 years	     (DTaP, IPV, Varicella, MMR) 
5 years	 
6, 8, and 10	 
11 years	  (HPV — in 2-3 doses given 6 months apart), Tdap, MCV) 
12 years	 
13 years	 
14 & 15 years	 
16 years	  (MCV booster) 
17 years	 

DTaP – Diphtheria, Tetanus, Pertussis. **Hep B** – Hepatitis B. **Hib** – Haemophilus influenza type b. **PCV** – Pneumococcal conjugate vaccine. **IPV** – Polio vaccine. **MMR** – Measles, Mumps, and Rubella. **HPV** – Human papillomavirus. **Tdap** – Diphtheria, Tetanus, Pertussis. **MCV** – Meningococcal conjugate vaccine.