

Vaccination Schedule for Children

Well-Child Check-up & Shots:
Children from birth to 17 years old



At your child's Well Visit talk with their doctor about what shots are right for them. If they are behind, your doctor can make a plan. Make sure to tell your doctor if your child will be traveling out of the U.S., they may need extra shots. Do not forget to talk about the flu shot.

Visit www.hpsj.com/well-visit to learn more.

 Well-Child Check-up	 Blood Screen	 Shots	 Vision Screen
 Hearing Screen	 Lead Screen	 Ask about the flu shot	
Age	Recommended Vaccines & Tests		
Birth	   (Hep B)		
3-5 days	 		
1 month	 		
2 months	   (DTaP, Hep B, Hib, PCV13, Rotavirus, IPV)		
4 months	  (DTaP, Hib, PCV13, Rotavirus, IPV)		
6 months	   (DTaP, Hep B, Lead screening, PCV13, Hib & Rotavirus if needed, IPV)		
9 Months			
12 months	    (Lead screening and anemia, MMR, Hep A, Varicella, Hib, PCV13)		
15-18 months	  (DTap)		
2 years	   (Lead screening, Hep A)		
3 years			
4 years	     (DTaP, IPV, Varicella, MMR)		
5 years			
6 – 10 years	  (Cholesterol screening between ages 9-11)		
11 years	   (HPV — in 2 doses given 5 months apart, or 3 doses if needed; Tdap, MCV)		
12 years			
13 years			
14 – 15 years			
16 years	  (MCV booster)		
17 years			

DTaP – Diphtheria, Tetanus, Pertussis. **Hep A/B** – Hepatitis A/B. **Hib** – Haemophilus influenza type b. **PCV13** – Pneumococcal conjugate vaccine. **IPV** – Polio vaccine. **MMR** – Measles, Mumps, and Rubella. **HPV** – Human papillomavirus. **Tdap** – Diphtheria, Tetanus, Pertussis. **MCV** – Meningococcal conjugate vaccine.